

ON MAXIMAL CURVES

RAINER FUHRMANN, ARNALDO GARCIA, AND FERNANDO TORRES

ABSTRACT. We study arithmetical and geometrical properties of *maximal curves*, that is, curves defined over the finite field $\mathbb{F}_{q^2}$ whose number of $\mathbb{F}_{q^2}$ -rational points reaches the Hasse-Weil upper bound. Under a hypothesis on non-gaps at a rational point, we prove that maximal curves are $\mathbb{F}_{q^2}$ -isomorphic to $y^q + y = x^m$, for some $m \in \mathbb{Z}^+$. As a consequence we show that a maximal curve of genus $g = (q-1)^2/4$ is $\mathbb{F}_{q^2}$ -isomorphic to the curve $y^q + y = x^{(q+1)/2}$.

0. INTRODUCTION

The interest on curves over finite fields was renewed after Goppa [Go] showed their applications to Coding Theory. One of the main features of linear codes arising from curves is the fact that one can state a lower bound for their minimum distance. This lower bound is meaningful only if the curve has many rational points. The subject of this paper is the study of *maximal curves*.

Let X be a projective, geometrically irreducible and non-singular algebraic curve defined over the finite field $\mathbb{F}_\ell$ with ℓ elements. A celebrated theorem of Weil states that:

$$\# X(\mathbb{F}_\ell) \leq \ell + 1 + 2g\sqrt{\ell},$$

where $X(\mathbb{F}_\ell)$ denotes the set of $\mathbb{F}_\ell$ -rational points of X and g is the genus of the curve. This bound was proved for elliptic curves by Hasse.

The curve X is called *maximal over $\mathbb{F}_\ell$* (in this case, ℓ must be a square; say $\ell = q^2$) if it attains the Hasse-Weil upper bound; that is,

$$\# X(\mathbb{F}_{q^2}) = q^2 + 1 + 2gq.$$

Ihara [Ih] shows that the genus of a maximal curve over $\mathbb{F}_{q^2}$ satisfies:

$$g \leq (q-1)q/2.$$

Rück and Stichtenoth [R-Sti] show that the Hermitian curve (that is, the curve given by $y^q + y = x^{q+1}$) is the unique (up to $\mathbb{F}_{q^2}$ -isomorphisms) maximal curve over $\mathbb{F}_{q^2}$ having genus $g = (q-1)q/2$.

It is also known that the genus of maximal curves over $\mathbb{F}_{q^2}$ satisfies (see [F-T] and the remark after Theorem 1.4 here):

$$g \leq (q-1)^2/4 \quad \text{or} \quad g = (q-1)q/2.$$

The Hermitian curve is a particular case of the following maximal curves over $\mathbb{F}_{q^2}$:

$$y^q + y = x^m, \text{ with } m \text{ being a divisor of } (q+1).$$

Note that the genus of the above curve is given by $g = (q-1)(m-1)/2$.

The paper was partially written while the first author was visiting the Instituto de Matemática Pura e Aplicada, Rio de Janeiro (Oct. 1995 - Jan. 1996) supported by CNPq and the third author was visiting ICTP (Trieste - Italy) supported by the International Atomic Energy Agency and UNESCO.

In Section 1 we derive properties of maximal curves. The main tools being the application to the linear system $\mathcal{D} = |(q+1)P_0|$, P_0 a rational point, of Stöhr-Voloch's approach [S-V] to the Hasse-Weil bound via Weierstrass Point Theory; and Castelnuovo's genus bound for curves in projective spaces: [C], [ACGH, p. 116], [Ra, Corollary 2.8]. A key result here is the fact that for any point P of the curve, the divisor $qP + \text{Fr}_X(P)$ is linearly equivalent to $\mathcal{D}$ (Corollary 1.2). This is a consequence of the particular fashion of the characteristic polynomial $h(t)$ of the Frobenius endomorphism of the Jacobian of the curve, that is, $h(t)$ is a power of a linear polynomial. This property also affects the geometry of the curve. More precisely, we show that maximal curves over $\mathbb{F}_{q^2}$ of genus $g \geq q-1$ are non-classical curves for the canonical morphism (Proposition 1.7). In some other cases one can deduce the non-classicality (for the canonical morphism) of the curve from the knowledge of $h(t)$. We will see this for the Deligne-Lusztig curve associated to the Suzuki group and to the Ree group (Proposition 1.8). The non-classicality of the curve corresponding to the Suzuki group was already proved in [G-Sti]. Our proof is different. It seems that the curve corresponding to the Ree group provides a new example of a non-classical curve.

In Section 2, we characterize the curves

$$y^q + y = x^m, \quad m \text{ being a divisor of } (q+1),$$

among the maximal curves over $\mathbb{F}_{q^2}$. This characterization being in terms of non-gaps at a rational point (Theorem 2.3). Finally in Section 3, applying the results of Section 2, we show that

$$y^q + y = x^{(q+1)/2}, \quad \text{with } q \text{ odd},$$

is the unique (up to $\mathbb{F}_{q^2}$ -isomorphisms) maximal curve over $\mathbb{F}_{q^2}$ with $g = (q-1)^2/4$.

1. MAXIMAL CURVES

Throughout this paper we use the following notation:

- By a curve we mean a projective, geometrically irreducible, non-singular algebraic curve defined over a finite field.
- Let k denote the finite field with q^2 elements, where q is a power of a prime p . Let $\bar{k}$ denote its algebraic closure.
- The symbol $X(k)$ (resp. $k(X)$) stands for the set of k -rational points (resp. for the field of k -rational functions) of a curve X defined over k .
- If $x \in k(X)$, then $\text{div}(x)$ (resp. $\text{div}_\infty(x)$) denotes the divisor (resp. the pole divisor) of the function x .
- Let P be a point of a curve. Then v_P (resp. $H(P)$) stands for the valuation (resp. for the Weierstrass non-gap semigroup) associated to P . We denote by $m_i(P)$ the i th non-gap at P .
- Let D be a divisor on X and $P \in X$. We denote by $\deg(D)$ the degree of D , by $\text{Supp}(D)$ the support of D , and by $v_P(D)$ the coefficient of P in D . If D is a k -divisor, we set

$$L(D) := \{f \in k(X) \mid \text{div}(f) + D \succeq 0\}, \quad \text{and} \quad \ell(D) := \dim_k L(D).$$

- The symbol " $\sim$ " denotes linear equivalence of divisors.
- The symbol g_d^r stands for a linear system of projective dimension r and degree d .

We first review some facts from Weierstrass Point Theory (see [Sch] and [S-V]).

Weierstrass points. Let X be a curve of genus g , and $\mathcal{D} = g_d^r$ be a base-point-free k -linear system on X . Then associated to a point $P \in X$ we have the Hermitian P -invariants $j_0(P) = 0 < j_1(P) < \dots < j_r(P) \leq d$ of $\mathcal{D}$ (also called the $(\mathcal{D}, P)$ -orders). This sequence is the same for all but finitely many points. These finitely many points P , where exceptional $(\mathcal{D}, P)$ -orders occur, are called the $\mathcal{D}$ -Weierstrass points of X . The Weierstrass points of the curve are those exceptional points obtained from the canonical linear system. A curve is called *non-classical* if the generic order sequence (for the canonical linear system) is different from $\{0, 1, \dots, g-1\}$.

Associated to the linear system $\mathcal{D}$ there exists a divisor R supporting exactly the $\mathcal{D}$ -Weierstrass points. Let $\epsilon_0 < \epsilon_1 < \dots < \epsilon_r$ denote the $(\mathcal{D}, Q)$ -orders for a generic point $Q \in X$. Then we have

$$(1.1) \quad \epsilon_i \leq j_i(P), \quad \text{for each } i = 0, 1, 2, \dots, r \text{ and for any point } P,$$

and also that

$$(1.2) \quad \deg(R) = (\epsilon_1 + \dots + \epsilon_r)(2g - 2) + (r + 1)d.$$

Associated to $\mathcal{D}$ we also have a divisor S whose support contains the set $X(k)$ of k -rational points on X . Its degree is given by

$$\deg(S) = (\nu_1 + \dots + \nu_{r-1})(2g - 2) + (q^2 + r)d,$$

where the ν_i 's form a subsequence of the ϵ_i 's. More precisely, there exists an integer I with $0 < I \leq r$ such that $\nu_i = \epsilon_i$ for $i < I$, and $\nu_i = \epsilon_{i+1}$ otherwise. Moreover, for $P \in X(k)$,

$$(1.3) \quad v_P(S) \geq \sum_{i=1}^r (j_i(P) - \nu_{i-1}), \quad \text{and} \quad \nu_i \leq j_{i+1}(P) - j_1(P), \quad \text{for each } i = 1, 2, \dots, r.$$

Maximal curves. We study some arithmetical and geometrical properties of maximal curves. To begin with we recall the following basic result concerning Jacobians. Let X be a curve, $\text{Fr}_{\mathcal{J}}$ the Frobenius endomorphism (relative to the base field) of the Jacobian $\mathcal{J}$ of X , and $h(t)$ the characteristic polynomial of $\text{Fr}_{\mathcal{J}}$. Let $h(t) = \prod_{i=1}^T h_i^{r_i}(t)$ be the factorization over $\mathbb{Z}[t]$ of $h(t)$. Then

$$(1.4) \quad \prod_{i=1}^T h_i(\text{Fr}_{\mathcal{J}}) = 0 \quad \text{on } \mathcal{J}.$$

This follows from the semisimplicity of $\text{Fr}_{\mathcal{J}}$ and the fact that the representation of endomorphisms of $\text{Fr}_{\mathcal{J}}$ on the Tate module is faithful (cf. [Ta, Thm. 2], [L, VI, §3]).

In the case of a maximal curve over $k = \mathbb{F}_{q^2}$, $h(t) = (t + q)^{2g}$. Therefore from (1.4) we obtain the following result, which is contained in the proof of [R-Sti, Lemma 1].

Lemma 1.1. *The Frobenius map $\text{Fr}_{\mathcal{J}}$ (relative to k) of the Jacobian $\mathcal{J}$ of a maximal curve over k acts as multiplication by $(-q)$ on $\mathcal{J}$.*

Let X be a maximal curve over k . Fix $P_0 \in X(k)$, and consider the map $f = f^{P_0} : X \rightarrow \mathcal{J}$ given by $P \rightarrow [P - P_0]$. We have

$$(1.5) \quad f \circ \text{Fr} = \text{Fr}_{\mathcal{J}} \circ f,$$

where Fr denotes the Frobenius morphism on X relative to k . Hence, from (1.5) and Lemma 1.1, we get:

Corollary 1.2. *For a maximal curve X over k , it holds*

$$\mathrm{Fr}(P) + qP \sim (q+1)P_0, \text{ for all points } P \text{ on } X.$$

It follows then immediately that

Corollary 1.3. ([R-Sti, Lemma 1]) *Let X be a maximal curve over k , $P_0, P_1 \in X(k)$. Then $(q+1)P_1 \sim (q+1)P_0$.*

Consider now the linear system $\mathcal{D} = g_{q+1}^{n+1} := |(q+1)P_0|$. Corollary 1.3 says that $\mathcal{D}$ is a k -invariant of the curve. In particular, its dimension $n+1$ is independent of the choice of $P_0 \in X(k)$. Moreover from Corollary 1.3 we have that $q+1 \in H(P_0)$; i.e., $(q+1)$ is a non-gap at a rational point, and hence $\mathcal{D}$ is base-point-free. From now on the letter $\mathcal{D}$ will always denote the linear system $|(q+1)P_0|$, P_0 a rational point, $(n+1)$ being its projective dimension, R will always mean the divisor supporting exactly the $\mathcal{D}$ -Weierstrass points, and Fr will always stand for the Frobenius morphism on X relative to k .

Theorem 1.4. *For a maximal curve X over k , the $\mathcal{D}$ -orders satisfy (notations being as above):*

- (i) $\epsilon_{n+1} = \nu_n = q$.
- (ii) $j_{n+1}(P) = q+1$ if $P \in X(k)$, and $j_{n+1}(P) = q$ otherwise; in particular, all rational points over k are $\mathcal{D}$ -Weierstrass points of X .
- (iii) $j_1(P) = 1$ for all points $P \in X$; in particular, $\epsilon_1 = 1$.
- (iv) If $n \geq 2$, then $\nu_1 = \epsilon_1 = 1$.

Proof. Statement (iii), for $P \in X(k)$, follows from (i), (ii) and the second inequality in (1.3). From Corollary 1.2 it follows the assertion (ii) and $\epsilon_{n+1} = q$. Furthermore, it also follows that $j_1(P) = 1$ for $P \notin X(k)$: in fact, let $P' \in X$ be such that $\mathrm{Fr}(P') = P$; then $P + qP' = \mathrm{Fr}(P') + qP' \sim (q+1)P_0$.

Now we are going to prove that $\nu_n = \epsilon_{n+1}$. Let $P \in X \setminus \{P_0\}$. Corollary 1.2 says that $\pi(\mathrm{Fr}(P))$ belongs to the osculating hyperplane at P , where π stands for the morphism associated to $\mathcal{D}$. This morphism π can be defined by a base $\{f_0, f_1, \dots, f_{n+1}\}$ of $L((q+1)P_0)$. Let x be a separating variable of $k(X) | k$. Then by [S-V, Prop. 1.4(c), Corollary 1.3] the rational function below is identically zero

$$w := \det \begin{pmatrix} f_0 \circ \mathrm{Fr} & \dots & f_{n+1} \circ \mathrm{Fr} \\ D_x^{\epsilon_0} f_0 & \dots & D_x^{\epsilon_0} f_{n+1} \\ \vdots & \vdots & \vdots \\ D_x^{\epsilon_n} f_0 & \dots & D_x^{\epsilon_n} f_{n+1} \end{pmatrix},$$

since it satisfies $w(P) = 0$ for a generic point P . Let I be the smallest integer such that the row $(f_0 \circ \mathrm{Fr}, \dots, f_{n+1} \circ \mathrm{Fr})$ is a linear combination of the vectors $(D_x^{\epsilon_i} f_0, \dots, D_x^{\epsilon_i} f_{n+1})$ with $i = 0, \dots, I$. Then according to [S-V, Prop. 2.1] we have

$$\{\nu_0 < \dots < \nu_n\} = \{\epsilon_0 < \dots < \epsilon_{I-1} < \epsilon_{I+1} < \dots < \epsilon_{n+1}\}.$$

That $\epsilon_1 = 1$ follows from statement (iii). Suppose that $\nu_1 > 1$. Since $j_1(P) = 1$ for all points of X , it follows from the proof of [H-V, Thm. 1] that

$$\# X(k) = (q+1)(q^2 - 1) - (2g - 2).$$

From the maximality of X , we then conclude $2g = (q-1) \cdot q$.

On the other hand, π is a birational morphism as follows from [Sti-X, Prop. 1] (see also Proposition 1.5(iv) here). Then Castelnuovo's genus bound for curves in projective spaces applied to the morphism π reads:

$$(1.6) \quad 2g \leq M \cdot (q - n + e) \leq \begin{cases} (2q - n)^2/4n, & \text{if } n \text{ is even} \\ ((2q - n)^2 - 1)/4n, & \text{if } n \text{ is odd,} \end{cases}$$

where M is the integer part of q/n and $e = q - M \cdot n$. We then conclude that $n = 1$ and this finishes the proof of the theorem. $\square$

Remark. For a maximal curve X with $n = 1$, we have $\nu_1 = \epsilon_2 = q > 1$. Then the proof above shows that $2g = (q - 1) \cdot q$. It then follows from [R-Sti] that the curve X is k -isomorphic to the Hermitian curve given by $y^q + y = x^{q+1}$. Also, if $n \geq 2$ then from Castelnuovo's formula (1.6) we get $g \leq (q - 1)^2/4$. This is the main result of [F-T].

The next proposition gives information on $\mathcal{D}$ -orders and non-gaps at points of X .

Proposition 1.5. *Let X be a maximal curve over k (notations being as before). Then:*

- (i) *For each point P on X , we have $\ell(qP) = n + 1$; i.e., we have the following behaviour for the non-gaps at P*

$$0 < m_1(P) < \cdots < m_n(P) \leq q < m_{n+1}(P).$$

- (ii) *If P is not rational over k , the numbers below are $\mathcal{D}$ -orders at the point P*

$$0 \leq q - m_n(P) < \cdots < q - m_1(P) < q.$$

- (iii) *If P is rational over k , the numbers below are exactly the $(\mathcal{D}, P)$ -orders*

$$0 < q + 1 - m_n(P) < \cdots < q + 1 - m_1(P) < q + 1.$$

In particular, if j is a $\mathcal{D}$ -order at a rational point P then $q + 1 - j$ is a non-gap at P .

- (iv) *If $P \in X(\mathbb{F}_{q^4}) \setminus X(k)$, then $q - 1$ is a non-gap at P . If $P \notin X(\mathbb{F}_{q^4})$, then q is a non-gap at P . If $P \in X(k)$, then q and $q + 1$ are non-gaps at P .*
- (v) *Let P be a non-Weierstrass point of X (for the canonical morphism) and suppose that $n \geq 2$, then we have for the non-gaps at P that $m_{n-1}(P) = q - 1$ and $m_n(P) = q$.*

Proof. Assertion (i) follows from Corollary 1.2. Let $m(P)$ be a non-gap at a point P of X with $m(P) \leq q$, then by definition there exists a positive divisor E disjoint from P with

$$E \sim m(P) \cdot P.$$

Summing up to both sides of the equivalence above the divisor $(q - m(P)) \cdot P + \text{Fr}(P)$, we get

$$E + (q - m(P)) \cdot P + \text{Fr}(P) \sim qP + \text{Fr}(P) \sim (q + 1)P_0.$$

This proves assertions (ii) and (iii). To prove assertion (iv) we just apply (as in [Har, IV, Ex. 2.6]) the Frobenius morphism to the equivalence in Corollary 1.2, getting

$$\text{Fr}^2(P) + (q - 1)\text{Fr}P \sim qP.$$

The fact that q and $q + 1$ are non-gaps at any rational point follows from assertion (iii) taking $j = 0$ and $j = 1$.

Now we are going to prove the last assertion (v). From assertion (iv) we know already

$$m_n(P) = q \quad \text{and} \quad m_{n-1}(P) \leq q - 1.$$

Suppose that $m_{n-1}(P) < (q-1)$. It then follows from Theorem 1.4 and the assertion (ii) above that the generic order sequence for the linear system $\mathcal{D}$ is as given below:

$$\epsilon_0 = 0 < \epsilon_1 = 1 < \epsilon_2 = q - m_{n-1}(P) < \cdots < \epsilon_n = q - m_1(P) < \epsilon_{n+1} = q.$$

On the other hand, we have that Equation (1.1) implies

$$m_i(Q) \leq m_i(P), \text{ for each } i \text{ and each } Q \in X.$$

Thus at a rational point $Q \in X$, it follows from assertion (iii) that:

$$v_Q(R) \geq \sum_{i=1}^{n+1} (j_i(Q) - \epsilon_i) = 1 + \sum_{i=1}^{n-1} (m_i(P) - m_i(Q) + 1) \geq n.$$

From the maximality of X , Equation (1.2) and [Ho, Thm. 1], we conclude that

$$n(q^2 + 2gq + 1) \leq \deg R \leq (n+2)\epsilon_{n+1}(g-1) + (n+2)(q+1).$$

Using that $\epsilon_{n+1} = q$, we finally have $nq^2 + qg(n-2) \leq 2$. This contradicts the assumption that $n \geq 2$. $\square$

Example 1.6. By Theorem 1.4 we have that all rational points of the curve are $\mathcal{D}$ -Weierstrass points. However, these sets may be different from each other as the following example shows:

Let X be the hyperelliptic curve defined by $x^2 + y^5 = 1$ over $\mathbb{F}_{81}$. The curve X is maximal because it is covered by the Hermitian curve $x^{10} + y^{10} = 1$ (see [Sti, Example VI.4.3]). It has genus 2 and at a generic point P , we have $m_7(P) = 9$. Hence we have $\mathcal{D} = |10P_0| = g_{10}^8$. All the canonical Weierstrass points are trivially rational points, and since $\#X(\mathbb{F}_{81}) = 118 > \#\{\text{Weierstrass points}\} = 6$, we have two possibilities for the $(\mathcal{D}, P)$ -orders at rational points, namely:

- (a) If P is a rational non-Weierstrass point; then its orders are 0, 1, 2, 3, 4, 5, 6, 7, 10.
- (b) If P is a Weierstrass point; then its orders are 0, 1, 2, 3, 4, 5, 6, 8, 10.

These computations follow from Proposition 1.5(iii). From the $\mathcal{D}$ -orders in (a) above, we conclude that the generic order sequence for $\mathcal{D}$ is 0, 1, 2, 3, 4, 5, 6, 7, 9. Hence, $\deg(R) = 164$ and $v_P(R) = 1$ (resp. $v_P(R) = 2$) if P satisfies (a) (resp. (b)) above. Since $\deg R - 112 \times 1 - 6 \times 2 = 40 > 0$, we then conclude that there exist non-rational $\mathcal{D}$ -Weierstrass points. The order sequence at such points must be 0, 1, 2, 3, 4, 5, 6, 8, 9 and so there exist 40 non-rational $\mathcal{D}$ -Weierstrass points, namely the fixed points of $\sigma \circ \text{Fr}$, where σ denotes the hyperelliptic involution.

By Proposition 1.5(v) we have that $q - n$ is a lower bound for the genus of a maximal curve over $\mathbb{F}_{q^2}$. We are going to show that classical (for the canonical morphism) maximal curves attain such a bound.

Proposition 1.7. *Let X be a maximal curve over $k = \mathbb{F}_{q^2}$ and let $g \geq 2$ be its genus. Then*

- (i) *If $g > q - n$ (with $n + 1 = \dim \mathcal{D}$ as before), then X is non-classical for the canonical morphism. In particular, this holds for $g \geq q - 1$.*
- (ii) *If X is hyperelliptic and the characteristic is two, then X has just one Weierstrass point for the canonical morphism.*

Proof. (i) If X is classical, then at a generic point P of the curve X we have

$$m_i(P) = g + i, \quad \forall i \in \mathbb{N}.$$

On the other hand, from Proposition 1.5(iv) we have $m_n(P) = q$. We then conclude that $g + n = q$. Now if $g \geq q - 1$ and X classical, then $n = 1$. Therefore from the remark after Theorem 1.4 we would have $2g = q(q - 1)$ and so $g = 1$, a contradiction.

(ii) Since X is hyperelliptic, the Weierstrass points are the fixed points of the hyperelliptic involution. Let P, Q be Weierstrass points of X (they exist because the genus is bigger than one). From $2P \sim 2Q$ and $2 \mid q$, we get $qP \sim qQ$. Therefore by Corollary 1.2,

$$qQ + \text{Fr}(Q) \sim qP + \text{Fr}(P) \sim qQ + \text{Fr}(P),$$

and so $\text{Fr}(P) \sim \text{Fr}(Q)$. This implies $\text{Fr}(P) = \text{Fr}(Q)$, and consequently $P = Q$. $\square$

Remark. Hyperelliptic maximal curves are examples of classical curves for the canonical morphism. It would be interesting to investigate the maximal curves that are both non-hyperelliptic and classical for the canonical morphism. Examples of such curves are the one of genus 3 over $\mathbb{F}_{25}$ listed by Serre in [Se, §4], and the generalizations of Serre's example obtained by Ibukiyama [I, Thm. 1].

Another question is whether or not the condition $g = q - n$ characterizes classical (for the canonical morphism) maximal curves.

Now we present two non-classical (for the canonical morphism) maximal curves over $\mathbb{F}_{q^2}$ of genus $g < q - 1$. These are the so-called Deligne-Lusztig curves associated to the Suzuki group and to the Ree group.

Proposition 1.8. (I) Let $s \in \mathbb{N}$, $r := 2^{2s+1}$, $r_0 := 2^s$, and consider the curve X over $\mathbb{F}_r$ defined by

$$y^r - y = x^{r_0}(x^r - x).$$

Then

- (i) ([H], [H-Sti], [Se]) The genus of X is $g = r_0(r - 1)$ and this curve is maximal over $\mathbb{F}_{r^4}$.
 - (ii) ([G-Sti]) The curve X is non-classical for the canonical morphism.
- (II) Let $s \in \mathbb{N}$, $r := 3^{2s+1}$, $r_0 := 3^s$, and consider the curve X over $\mathbb{F}_r$ defined by

$$y^r - y = x^{r_0}(x^r - x), \quad z^r - z = x^{2r_0}(x^r - x).$$

Then

- (i) ([H], [P], [Se]) The genus of X is $g = 3r_0(r - 1)(r + r_0 + 1)$ and this curve is maximal over $\mathbb{F}_{r^6}$.
- (ii) The curve X is non-classical for the canonical morphism.

Proof. We first set some notations. We write Fr for the Frobenius morphism on X relative to $\mathbb{F}_r$ and $h_i(t)$ for the characteristic polynomial of the Frobenius endomorphism (relative to $\mathbb{F}_{r^i}$) of the Jacobian of X .

(I) From [H, Prop. 4.3], [H-Sti], [Se] we know that $g = r_0(r - 1)$ and that $h_1(t) = (t^2 + 2r_0t + r)^g$. If a_1 and a_2 denote the roots of $h_1(t)$, then we have $a_1 + a_2 = -2r_0$ and $a_1a_2 = r$. It then follows easily that $(a_1a_2)^4 = r^4$ and $a_1^4 + a_2^4 = -2r^2$, and hence that $h_4(t) = (t + r^2)^{2g}$. This shows the maximality over $\mathbb{F}_{r^4}$.

Now by (1.4) we have $\text{Fr}_{\mathcal{J}}^2 + 2r_0\text{Fr}_{\mathcal{J}} + r = 0$ on $\mathcal{J}$ and then by (1.5) we obtain

$$\text{Fr}^2(P) + 2r_0\text{Fr}(P) + rP \sim (1 + 2r_0 + r)P_0,$$

for all $P \in X$, $P_0 \in X(\mathbb{F}_r)$. Now applying Fr to the equivalence above we get

$$\text{Fr}^3(P) + (2r_0 - 1)\text{Fr}^2(P) + (r - 2r_0)\text{Fr}(P) \sim rP.$$

Hence we conclude that $r \in H(P)$ at a general point $P \in X$, and since $g \geq r$, that X is non-classical for the canonical morphism.

(II) From [H, Prop. 5.3], [P], [Se] we already know the formula for the genus and that $h_1(t) = (t^2 + r)^a(t^2 + 3r_0t + r)^b$ with $a, b \in \mathbb{N}$ and $a + b = g$. Let a_1, a_2 denote the roots of some factor of $h_1(t)$. Then in either case we get $(a_1a_2)^6 = r^6$ and $a_1^6 + a_2^6 = -2r^3$ and hence that $h_6(t) = (t + r^3)^{2g}$. This shows the maximality of X over $\mathbb{F}_{r^6}$.

Finally as in the proof of item (I) we conclude that $r^2 \in H(P)$ for a generic $P \in X$. Since $g \geq r^2$ the assertion follows. $\square$

To finish this section on maximal curves, we study some properties involving the morphism $\pi : X \rightarrow \mathbb{P}^{n+1}$ associated to the linear system $\mathcal{D} = |(q+1)P_0|$.

Proposition 1.9. *The following statements are equivalent:*

- (i) *The morphism π is a closed embedding, i.e. X is k -isomorphic to $\pi(X)$.*
- (ii) *For all $P \in X(\mathbb{F}_{q^4})$, we have that $\pi(P) \in \mathbb{P}^{n+1}(k) \Leftrightarrow P \in X(k)$.*
- (iii) *For all $P \in X(\mathbb{F}_{q^4})$, we have that q is a non-gap at P .*

Proof. Let $P \in X$. Since $j_1(P) = 1$ (cf. Theorem 1.4(iii)), we have that $\pi(X)$ is non-singular at all branches centered at $\pi(P)$. Thus π is an embedding if and only if π is injective.

Claim. *We have $\pi^{-1}(\pi(P)) \subseteq \{P, \text{Fr}(P)\}$ and if π is not injective at the point P , then $P \in X(\mathbb{F}_{q^4}) \setminus X(k)$ and $\pi(P) \in \mathbb{P}^{n+1}(k)$.*

From Corollary 1.2 it follows that $\pi^{-1}(\pi(P)) \subseteq \{P, \text{Fr}(P)\}$. Now if π is not injective at P , then $P \notin X(k)$ and, since $P \in \pi^{-1}(\pi(\text{Fr}(P))) \subseteq \{\text{Fr}(P), \text{Fr}^2(P)\}$, we have $\text{Fr}^2(P) = P$, i.e. $P \in X(\mathbb{F}_{q^4}) \setminus X(k)$. Furthermore we have $\pi(P) = \pi(\text{Fr}(P)) = \text{Fr}(\pi(P))$, i.e. $\pi(P) \in \mathbb{P}^{n+1}(k)$. This proves the claim above.

From this claim the equivalence (i) $\Leftrightarrow$ (ii) follows immediately. As to the implication (i) $\Rightarrow$ (iii), we know that $\dim |\text{Fr}(P) + qP - P - \text{Fr}(P)| = \dim |\text{Fr}(P) + qP| - 2$ (Corollary 1.2 and [Har, Prop. 3.1(b)]), i.e. $\ell((q-1)P) = n$ and so $q \in H(P)$, by Proposition 1.5(i).

Finally we want to conclude from (iii) that π is an embedding. According to the above claim it is sufficient to show that $\pi^{-1}(\pi(P)) = \{P\}$, for $P \in X(\mathbb{F}_{q^4}) \setminus X(k)$. Let then $P \in X(\mathbb{F}_{q^4})$. Since we have $q \in H(P)$, there is a divisor $D \in |qP|$ with $P \notin \text{Supp}(D)$. In particular,

$$\text{Fr}(P) + D \sim \text{Fr}(P) + qP \sim (q+1)P_0,$$

and then $\pi^{-1}(\pi(\text{Fr}(P))) \subseteq \text{Supp}(\text{Fr}(P) + D)$. So if $\pi^{-1}(\pi(P)) = \{P, \text{Fr}(P)\}$, then we would have that $P \in \text{Supp}(D)$, a contradiction. This means altogether that π is injective and so indeed a closed embedding. $\square$

Remark. Condition (iii) above is satisfied whenever $q \geq 2g$, and in most of the well known examples of maximal curves the morphism π is always an embedding. Then a natural question is whether or not π is an embedding for an arbitrary maximal curve. We conjecture that this property is a necessary condition for a maximal curve being covered by the Hermitian curve.

Proposition 1.10. *Suppose that $\pi : X \rightarrow \mathbb{P}^{n+1}$ is a closed embedding. Let $P_0 \in X(k)$ and assume furthermore that there exist $r, s \in H(P_0)$ such that all non-gaps at P_0 less than or equal to $q + 1$ are generated by r and s . Then the semigroup $H(P_0)$ is generated by r and s . In particular, the genus of X is equal to $(r - 1)(s - 1)/2$.*

Proof. Let $x, y \in k(X)$ with $\text{div}_\infty(x) = sP_0$ and $\text{div}_\infty(y) = rP_0$. Since we have that $q, q + 1 \in H(P_0)$, then the numbers r and s are coprime. Let $\pi_2 : X \rightarrow \mathbb{P}^2$, be given by $P \mapsto (1 : x(P) : y(P))$. Then the curves X and $\pi_2(X)$ are birational and the image $\pi_2(X)$ is a plane curve given by an equation of the type below:

$$x^r + \beta y^s + \sum_{is+jr < rs} \alpha_{ij} x^i y^j = 0,$$

where $\beta, \alpha_{ij} \in k$ and $\beta \neq 0$. We are going to prove that $\pi_2(P)$ is a non-singular point of the curve $\pi_2(X)$ for all $P \neq P_0$. From this it follows that $g = (r - 1)(s - 1)/2$ and also that $H(P_0) = \langle r, s \rangle$ (see [Ful, Ch. 7], [To]).

Let $1, f_1, \dots, f_{n+1}$ be a basis of $L((q + 1)P_0)$, where $n + 1 := \dim |(q + 1)P_0|$. Then there exist polynomials $F_i(T_1, T_2) \in k[T_1, T_2]$ for $i = 1, \dots, n + 1$, such that

$$f_i = F_i(x, y), \quad \text{for } i = 1, \dots, n + 1.$$

The existence of these polynomials follows from the hypothesis on the non-gaps at P_0 less than or equal to $(q + 1)$.

Consider the maps $\pi|_{(X \setminus \{P_0\})} : X \setminus \{P_0\} \rightarrow \mathbb{A}^{n+1}$ given by $P \mapsto (f_1(P), \dots, f_{n+1}(P))$; $\pi_2|_{(X \setminus \{P_0\})} : X \setminus \{P_0\} \rightarrow \mathbb{A}^2$, $P \mapsto (x(P), y(P))$; and $\phi : \mathbb{A}^2 \rightarrow \mathbb{A}^{n+1}$, given by $(p_1, p_2) \mapsto (F_1(p_1, p_2), \dots, F_{n+1}(p_1, p_2))$. Then the following diagram is commutative

$$\begin{array}{ccc} & X \setminus \{P_0\} & \\ \pi_2 \swarrow & & \searrow \pi \\ \mathbb{A}^2 & \xrightarrow{\phi} & \mathbb{A}^{n+1} \end{array}$$

Thus we have for a point P of $X \setminus \{P_0\}$ and the corresponding local rings assigned to $\pi(P), \pi_2(P)$ the commutative diagram

$$\begin{array}{ccc} & O_{\pi(X), \pi(P)} & \\ f \swarrow & & \searrow c \\ O_{\pi_2(X), \pi_2(P)} & \xrightarrow{h} & O_{X, P} \end{array}$$

where h is injective since $k(X) = k(x, y)$, and c is an isomorphism by assumption. Thus $\pi_2 X$ is non-singular at $\pi_2 P$. $\square$

2. CERTAIN MAXIMAL CURVES

The curves we have in mind in this section are the ones given by (see [G-V] and [Sch]):

$$y^q + y = x^m, \quad \text{with } m \text{ being a divisor of } (q + 1).$$

These are maximal curves (with $2g = (m-1)(q-1)$) since they are covered by the Hermitian curve. If P_0 is the unique point at infinity of this curve, then the semigroup of non-gaps at P_0 is generated by m and q and we have:

$$m \cdot n = q + 1, \quad \text{where} \quad (n+2) = \ell((q+1)P_0). \quad (**)$$

The goal of this section is to give a proof that the above condition $(**)$ on non-gaps at a rational point P_0 characterizes the curves $y^q + y = x^m$ among the maximal curves over the finite field k .

Proposition 2.1. *Let X be a maximal curve of genus g . Suppose that there exists a rational point $P_0 \in X(k)$ such that $n \cdot m = q + 1$, with m being a non-gap at P_0 . Then, we have $2g = (q-1)(m-1)$. Also, there are at most two types of $(\mathcal{D}, P)$ -orders at rational points $P \in X(k)$:*

Type 1. *The $\mathcal{D}$ -orders at P are $0, 1, 2, 3, \dots, n, q+1$. In this case we have $v_P(R) = 1$.*

Type 2. *The $\mathcal{D}$ -orders at P are $0, 1, m, 2m, \dots, (n-1)m, q+1$. In this case we have $w_2 := v_P(R) = n((n-1)m - n - 1)/2 + 2$.*

Moreover, the set of $\mathcal{D}$ -Weierstrass points of X coincides with the set of its k -rational points, and the order sequence for $\mathcal{D}$ is $0, 1, 2, \dots, n, q$.

Proof. The morphism π can be defined by $(1 : y : \dots : y^{n-1} : x : y^n)$, where $x, y \in k(X)$ are functions such that

$$\operatorname{div}_\infty(x) = qP_0 \quad \text{and} \quad \operatorname{div}_\infty(y) = mP_0.$$

The set of $\mathcal{D}$ -orders at P_0 is of Type 2, as follows from Proposition 1.5(iii).

Let $P \in X \setminus \{P_0\}$. From the proof of [S-V, Thm. 1.1] and letting $z = y - y(P)$, we have

$$(2.1) \quad v_P(z), \dots, nv_P(z)$$

are $(\mathcal{D}, P)$ -orders. Thus, considering a non-ramified point for $y : X \rightarrow \mathbb{P}^1$, we conclude that the order sequence of the linear system $\mathcal{D}$ is given by

$$\epsilon_i = i \quad \text{for } i = 1, \dots, n, \quad \text{and} \quad \epsilon_{n+1} = q.$$

If P is a rational point, by Theorem 1.4, we know that 1 and $(q+1)$ are $(\mathcal{D}, P)$ -orders. We consider two cases:

- (1) $v_P(z) = 1$: This implies that the point P is of Type 1.
- (2) $v_P(z) > 1$: From assertion (2.2) above, it follows $n \cdot v_P(z) = q + 1$ and hence $v_P(z) = m$. Then, we have that the point P is of Type 2.

If P is not a rational point, by Theorem 1.4, we have that $j_{n+1}(P) = q$. If $v_P(z) > 1$ and using assertion (2.1), we get

$$n \cdot v_P(z) = q = n \cdot m - 1.$$

Hence $n = 1$ and the $(\mathcal{D}, P)$ -orders are $0, 1, q$. This shows that P is not a $\mathcal{D}$ -Weierstrass point. If $v_P(z) = 1$, again from assertion (2.1), we have that

$$0, 1, 2, \dots, n, q$$

are the $\mathcal{D}$ -orders at the point P ; i.e., P is not a $\mathcal{D}$ -Weierstrass point. This shows the equality of the two sets below:

$$\{\mathcal{D} - \text{Weierstrass points of } X\} = \{k - \text{rational points of } X\}.$$

The assertions on $v_P(R)$ follow from [S-V, Thm. 1.5].

Let T_1 (resp. T_2) denote the number of rational points $P \in X(k)$ whose $(\mathcal{D}, P)$ -orders are of Type 1 (resp. Type 2). Thus we have from the equality in (1.2)

$$\deg(R) = (n(n+1)/2 + q)(2g-2) + (n+2)(q+1) = T_1 + w_2 T_2.$$

Riemann-Hurwitz applied to $y : X \rightarrow \mathbb{P}^1(\bar{k})$ gives

$$2g-2 = -2m + (m-1)T_2.$$

Since $T_1 + T_2 = \#X(k) = q^2 + 2gq + 1$, and using the two equations above, we conclude after tedious computations that $2g = (m-1)(q-1)$. This finishes the proof of the proposition. $\square$

Now we are going to prove that maximal curves as in Proposition 2.1 are isomorphic to $y^q + y = x^m$. To begin with we first generalize [R-Sti, Lemma 5].

Lemma 2.2. *Notations and hypotheses as above. Then, the extension $k(X) \mid k(y)$ is a Galois cyclic extension of degree m .*

Proof. From the proof of Proposition 2.1 we see that the extension $k(X) \mid k(y)$ is ramified exactly at the rational points of Type 2 and that $T_2 = (q+1)$. Moreover, this extension is totally ramified at those points. Viewing the function y on X as a morphism of degree m

$$y : X \rightarrow \mathbb{P}^1(\bar{k}),$$

we consider the elements of the finite field k belonging to the set below

$$V_1 = \{y(P) \mid P \in X(k) \text{ is of Type 1}\}.$$

Since $T_2 = (q+1)$ and the rational points of Type 2 are totally ramified for the morphism y , we must have $\#V_1 \leq (q^2 - q)$. Now, above each element of the set V_1 there are at most m rational points of the curve X , those points being necessarily of Type 1, and hence:

$$\#X(k) = q^2 + 1 + 2gq = T_1 + T_2 \leq m \cdot (q^2 - q) + (q+1).$$

From the genus formula in Proposition 2.1, we then conclude that $\#V_1 = (q^2 - q)$ and also that above each element of the set V_1 there are exactly m rational points of the curve.

Now the proof continues as in the proof of [R-Sti, Lemma 5]. We will repeat their argument here for completeness. Let $\tilde{F}$ be the Galois closure of the extension $k(X) \mid k(y)$. The field k is still algebraically closed in $\tilde{F}$ since the elements of the set V_1 split completely in $k(X) \mid k(y)$. Moreover the extension $\tilde{F} \mid k(X)$ is unramified, as follows from Abhyankar's lemma [Sti, ch.III.8]. Hence,

$$2\tilde{g} - 2 = [\tilde{F} : F](2g - 2),$$

where $\tilde{g}$ denotes the genus of the field $\tilde{F}$. The $(q^2 - q)$ elements of the set V_1 split completely in $\tilde{F}$ and then they give rise to $(q^2 - q)m[\tilde{F} : F]$ rational points of $\tilde{F}$ over k . Then, from the Hasse-Weil bound, we conclude

$$(q^2 - q)m[\tilde{F} : F] \leq q^2 + 2q + (2\tilde{g} - 2)q = q^2 + 2q + [\tilde{F} : F](2g - 2)q.$$

Substituting $2g = (m-1)(q-1)$ in the inequality above, we finally get:

$$[\tilde{F} : F] \leq \frac{q+2}{q+1} \text{ and hence } \tilde{F} = F.$$

Note that the extension is cyclic since there exist rational points (those of Type 2) that are totally ramified for the morphism y . $\square$

Theorem 2.3. *Let X be a maximal curve of genus g such that there exists a rational point $P_0 \in X(k)$ with $m \cdot n = (q + 1)$, where m is a non-gap at P_0 . Then the curve X is k -isomorphic to the curve given by the equation:*

$$y_1^q + y_1 = x_1^m.$$

Proof. We know that $k(X) \mid k(y)$ is a Galois cyclic extension of degree m and moreover that the functions $1, y, y^2, \dots, y^{n-1}$ and x form a basis for $L(qP_0)$. Let σ be a generator of the Galois group of $k(X) \mid k(y)$. Since P_0 is totally ramified, then $\sigma(P_0) = P_0$ and hence $\sigma(L(qP_0)) = L(qP_0)$. Note that the functions $1, y, y^2, \dots, y^{n-1}$ form a basis for the subspace $L((n-1)mP_0)$ and that σ acts as the identity on this subspace. Since m and q are relatively prime, we can diagonalize σ on $L(qP_0)$. Take then a function $v \in L(qP_0)$, $v \notin L((n-1)mP_0)$, satisfying $\sigma(v) = \lambda v$, with λ a primitive m -th root of 1.

Then denoting by N the norm of $k(X) \mid k(y)$, we get $N(v) = (-1)^{m+1} \cdot v^m$.

Hence $v^m \in k(y)$ and since it has poles only at P_0 , we must have $v^m = f(y) \in k[y]$. Since $\text{div}_\infty(v) = \text{div}_\infty(x) = qP_0$, we see that $\deg f(y) = q$. Now from the fact that there are exactly $(q+1)$ totally ramified points of $k(X) \mid k(y)$ and that all of them are rational, we conclude that $f(y) \in k[y]$ is separable and has all its roots in k . After a k -rational change of coordinates, we may assume that $f(0) = 0$. Then, we get the following description for the set V_1 : $V_1 = \{\alpha \in k \mid f(\alpha) \neq 0\}$. Knowing that all points of X above V_1 are rational points over k and from the equation $v^m = f(y)$, we get:

$$f^n(y) \equiv f^{nq}(y) \pmod{y^{q^2} - y}. \quad (*)$$

Claim. $f(y) = a_1 y + a_q y^q$, with $a_1, a_q \in k^*$.

We set $f(y) = \sum_{i=1}^q a_i y^i$ and $f^n(y) = \sum_{i=n}^{nq} b_i y^i$. Clearly, the fact that $a_1, a_q \in k^*$ follows from the fact that $f(y)$ is separable of degree q . Suppose that the set I below is non-empty

$$I = \{2 \leq i \leq q-1 \mid a_i \neq 0\},$$

and then define

$$t = \min I \quad \text{and} \quad j = \max I.$$

Clearly, we have $b_{(n-1)q+j} = n \cdot a_q^{n-1} \cdot a_j \neq 0$. Since the unique solution for i in the congruence $i q \equiv (n-1)q + j \pmod{q^2 - 1}$, i being smaller than q^2 , is the one given by $i = (n-1) + j q$, it follows from (*) above that $b_{(n-1)q+j} = b_{(n-1)+jq}^q \neq 0$.

It now follows that $\deg(f^n(y)) = nq \geq (n-1) + jq$ and hence we get that $n-j \geq 1$ if $n \geq 2$. Note that if $n = 1$, then we get $j \leq 1$ and the proof of the claim is complete in this case. From now on we then assume $n \geq 2$. We then conclude that $t \leq j \leq (n-1)$. Note that then $(n-1) + t < q$.

Clearly, we also have $b_{n-1+t} = n \cdot a_1^{n-1} \cdot a_t \neq 0$. Since the unique solution for i in the congruence $i q \equiv n-1+t \pmod{q^2 - 1}$, i being smaller than q^2 , is the one given by $i = (n-1+t)q$, it follows from (*) above that $b_{n-1+t} = b_{(n-1+t)q}^q \neq 0$.

As before, it now follows that $nq \geq (n-1+t)q$, and hence $t \leq 1$. This gives the desired contradiction and hence the set I is empty, thereby proving the claim.

Now we are in a position to finish the proof of the theorem. Denoting

$$f(k) = \{f(\alpha) \mid \alpha \in k\} \quad \text{and} \quad H = \{\beta^m \mid \beta \in k\},$$

we have that $\mathbb{F}_q^*$ is a subgroup of $H \setminus \{0\}$ of index equal to n . Moreover, using the fact that above V_1 there are only rational points, we have:

$$f(k) \subseteq H = \bigcup_{\ell=0}^{n-1} \xi^{\ell \cdot m} \mathbb{F}_q,$$

where ξ denotes a primitive element of the field k ; i.e., ξ is a generator for the multiplicative cyclic group k^* . Since $f(k)$ is a $\mathbb{F}_q$ -linear subspace of k as follows from the above claim, we conclude that its dimension is one and hence that $f(k) = \xi^{r \cdot m} \mathbb{F}_q$, for some r . Finally, putting $x_1 = \xi^{-r} v$ and $y_1 = \epsilon y$, where ϵ is the unique element of k^* satisfying

$$\text{Tr}(\epsilon \alpha) = \xi^{-r \cdot m} f(\alpha), \quad \forall \alpha \in k,$$

we conclude the proof of the theorem (Tr being the trace operator in $k(X) \mid k(y)$). $\square$

Remark. Notations being as above. Suppose that $m \cdot n \leq q + 1$ (m being a non-gap at some rational point P_0 of X). Then, we have $q + 1 \geq m \cdot n \geq m_n(P_0) = q$, where the last equality follows from Proposition 1.5(iv). In case that $m \cdot n = q$, we conjecture that $2g = (m - 1)q$ and the curve is $\mathbb{F}_{q^2}$ -isomorphic to a curve given by

$$F(y) = x^{q+1},$$

where $F(y)$ is a $\mathbb{F}_p$ -linear polynomial of degree m . We have not been able to prove this possible result yet.

We notice that if one could show that the morphism $\pi : X \rightarrow \mathbb{P}^{n+1}$ is a closed embedding, then by Proposition 1.10 we would have the claimed formula for g .

Finally we also notice that $(m_1(P) - 1)q/2$, $P \in X(k)$, is an upper bound for the genus of maximal curves. This follows from [Le, Thm. 1(b)].

Example 2.4. There exist maximal curves that do not satisfy the hypothesis of Theorem 2.3. We give two such examples below:

(i) Let X be the maximal curve over $\mathbb{F}_{25}$ and genus $g = 3$ listed by Serre in [Se, §4]. Let $m, 5, 6$ be the first three non-gaps at $P \in X(\mathbb{F}_{25})$. Here we have $6P_0 = g_6^3$. We claim that $m = 4$ (and so $nm > q + 1$). Indeed, if $m = 3$ by Proposition 2.1 we would have $g = 4$.

This example also shows a maximal curve where all the rational points are non-Weierstrass points: in fact, since $5 = \text{char}(k) > 2g - 2$ the curve is classical.

(ii) Let X be a maximal curve over $\mathbb{F}_{q^2}$ of genus g . Suppose that $q \geq 2g + 2$ (e.g. the maximal curves in Proposition 1.8 here, [G-VI, Thm. 3.12, Thm. 3.16], [I, Thm. 1]). Then X does not satisfy the hypothesis of Theorem 2.3. In fact, for $P_0 \in X(k)$ we have $m_{q+i}(P_0) = 2g + i$ and then $n = q - g$. Therefore $m_1(P_0)n \geq 2n \geq q + 2$, the last inequality following from $q \geq 2g + 2$.

3. MAXIMAL CURVES OF GENUS $(q - 1)^2/4$.

As an interesting application of the preceding section we prove:

Theorem 3.1. *Let X be a maximal curve over $\mathbb{F}_{q^2}$ of genus $g = (q - 1)^2/4$. Then the curve X is $\mathbb{F}_{q^2}$ -isomorphic to the one given by*

$$y^q + y = x^{q+1/2}.$$

Proof. From Equation (1.6) which is Castelnuovo's genus bound applied to the linear system $|(q+1)P_0|$, we have $n = 2$ (see remark before Proposition 1.5). From Theorem 2.3, it suffices to prove the existence of a rational point P over k with $m_1(P) = (q+1)/2$. This is clearly true for $q = 3$ (since $g = 1$ in this case) and hence we can assume $q \geq 5$.

We prove firstly some lemmas:

Lemma 3.2. *Let P be a rational point over k of the curve X (hypothesis being as in Theorem 3.1). Then we have that $\ell((2q+2)P) = 9$ and that either $m_1(P) = (q-1)$ or $m_1(P) = (q+1)/2$. Moreover, the divisor $(2g-2)P$ is a canonical divisor.*

Proof. Let $m_i = m_i(P)$ be the i -th non-gap at the rational point P . We have the following list of non-gaps at P :

$$0 < m_1 < m_2 = q < m_3 = q+1 \leq 2m_1 < m_1 + m_2 < m_1 + m_3 \leq 2m_2 < m_2 + m_3 < 2m_3.$$

The inequality $2m_1 \geq (q+1)$ follows from the fact that $n = 2$ and q odd. Clearly,

$$\begin{aligned} m_3 = 2m_1 & \quad \text{if and only if} \quad m_1 = (q+1)/2; \quad \text{and} \\ m_1 + m_3 = 2m_2 & \quad \text{if and only if} \quad m_1 = (q-1). \end{aligned}$$

Since $q \geq 5$, one cannot have both equalities above simultaneously. From the above list of non-gaps at P , it then follows that $\ell((2q+2)P) \geq 9$. Moreover, after showing that $\ell((2q+2)P) = 9$, it also follows that either $m_1 = (q+1)/2$ or $m_1 = (q-1)$. Let $\pi_2: X \rightarrow \mathbb{P}^{r+1}$ be the morphism associated to the linear system $|(2q+2)P|$; we already know that $r \geq 7$ and we have to show that $r = 7$. Castelnuovo's bound for the morphism π_2 gives

$$2g = \frac{(q-1)^2}{2} \leq M \cdot (d-1-(r-e)), \quad (*)$$

where $d = 2q+2$, $M = \left\lfloor \frac{d-1}{r} \right\rfloor$ and $d-1 = M \cdot r + e$. Since $(r-e) \geq 1$ we have $d-1-(r-e) \leq 2q$, and hence

$$(q-1)^2 \leq 4qM \quad \text{and then} \quad q^2 - q \leq 4qM,$$

since the right hand side above is a multiple of q . For $r \geq 9$, we now see that

$$q-1 \leq 4M \leq 4 \cdot \frac{2q+1}{9}, \quad \text{and then} \quad q \leq 13.$$

The cases $q \leq 13$ are discarded by direct computations in Equation (*) above, and hence we have $r \leq 8$. Now we use again Equation (*) to discard also the possibility $r = 8$. Since q is odd, we have

$$2q+1 \equiv 3 \pmod{8} \quad \text{or} \quad 2q+1 \equiv 7 \pmod{8}.$$

It then follows

$$\begin{cases} M = (q-1)/4 \\ \text{and } e = 3 \end{cases} \quad \text{or} \quad \begin{cases} M = (q-3)/4 \\ \text{and } e = 7. \end{cases}$$

Substituting these two possibilities above in Equation (*), one finally gets the desired contradiction; i.e., one gets

$$(q-1)^2 \leq (q-1)(q-2) \quad \text{or} \quad (q-1)^2 \leq (q-3) \cdot q.$$

Now we prove the last assertion of the lemma. One can easily check that both semigroups H_1 and H_2 below are symmetric, with exactly $g = (q - 1)^2/4$ gaps:

$$H_1 = \langle (q - 1), q, q + 1 \rangle \quad \text{and} \quad H_2 = \langle \frac{q + 1}{2}, q \rangle.$$

At a rational point P on X the Weierstrass semigroup $H(P)$ must then be equal to H_1 or H_2 . Hence the semigroup $H(P)$ is necessarily symmetric and the last assertion follows. $\square$

Lemma 3.3. *Let $\mathcal{D} = |(q + 1)P|$ with P being a rational point of X (hypothesis as in Theorem 3.1). Then at any non-rational point Q of X , the $(\mathcal{D}, Q)$ -orders are $0, 1, 2, q$. In particular the order sequence for $\mathcal{D}$ is $0, 1, 2, q$, and the set of $\mathcal{D}$ -Weierstrass points is exactly the set of rational points.*

Proof. Let $0, 1, j, q$ be the $(\mathcal{D}, Q)$ -orders. Consider the following set S :

$$S = \{0, 1, 2, j, j + 1, 2j, q, q + 1, q + j, 2q\}.$$

The set S consists of $(2\mathcal{D}, Q)$ -orders, and hence from Lemma 3.2 we must have $\# S \leq 9$. This eliminates the possibilities

$$3 \leq j \leq (q - 1)/2 \quad \text{and} \quad \frac{q + 3}{2} \leq j \leq q - 2,$$

and it then follows that $j \in \{2, (q + 1)/2, q - 1\}$. From Lemma 3.2 we know that

$$(2q - 2)P = \frac{(q + 1)(q - 3)}{2}P \quad \text{is canonical.}$$

Then the following set $S(j)$ consists of orders at Q for the canonical morphism

$$S(j) = \{a + bj + cq \mid a, b, c \in \mathbb{N} \text{ with } a + b + c \leq \frac{q - 3}{2}\}.$$

One can check that $\# S(j) = (q - 1)^2/4$ if the value of j belongs to $\{2, (q + 1)/2, q - 1\}$, and hence that $S(j)$ consists of all canonical orders at the point Q . Then the set $H(j)$ below is necessarily a semigroup:

$$H(j) = \mathbb{N} \setminus (1 + S(j)).$$

This semigroup property on $H(j)$ is only satisfied for the value $j = 2$, as one checks quite easily, and this finishes the proof of this lemma. $\square$

Now we turn back to the proof of Theorem 3.1. Suppose that $m_1(P) = (q - 1)$ at all rational points P on the curve. It then follows from Proposition 1.5(iii) that the $(\mathcal{D}, P)$ -orders are $0, 1, 2, q + 1$ and hence $v_P(R) = 1$, where R is the divisor supporting the $\mathcal{D}$ -Weierstrass points. On the other hand, we have

$$\deg R - \# X(k) = 3(2g - 2) - (q - 3)(q + 1) = \frac{1}{2}(q + 1)(q - 3).$$

Since $q \geq 5$ and $v_P(R) = 1$ for P rational, we would then conclude the existence of non-rational points that are $\mathcal{D}$ -Weierstrass points. This contradicts Lemma 3.3 and hence, from Lemma 3.2, we finally conclude the existence of a rational point P satisfying

$$m_1(P) = (q + 1)/2.$$

$\square$

We can explore further the idea of the above proofs to obtain a partial analogue of the main result of [F-T], namely

Scholium 3.4. *Let X be a maximal curve over k whose genus g satisfies*

$$(q^2 - 3q + 2)/4 < g \leq (q - 1)^2/4.$$

If q is odd, neither q is a power of 3 nor $q \not\equiv 3 \pmod{4}$, then $g = (q - 1)^2/4$.

Notice that Example 2.4(i) shows that the hypothesis on g above is sharp. This Scholium is the first step toward a characterization of a maximal curve whose genus is $\frac{q-1}{2}(\frac{q+1}{t} - 1)$ with $t \geq 3$.

REFERENCES

- [ACGH] Arbarello, E.; Cornalba, M.; Griffiths, P.A. and Harris, J.: Geometry of algebraic curves, Vol. I, Springer-Verlag, New York 1985.
- [C] Castelnuovo, G.: Ricerche di geometria sulle curve algebriche, Atti. R. Acad. Sci. Torino **24**, 196–223 (1889).
- [F-T] Fuhrmann, R.; Torres, F.: The genus of curves over finite fields with many rational points, Manuscripta Math. **89**, 103–106 (1996).
- [Ful] Fulton, W.: Algebraic curves. An introduction to algebraic curves (second edition). W.A. Benjamin, INC 1974.
- [G-Sti] Garcia, A.; Stichtenoth, H.: Elementary abelian p -extensions of algebraic function fields, Manuscripta Math. **72**, 67–79 (1991).
- [G-V] Garcia, A.; Viana, P.: Weierstrass points on certain non-classical curves, Arch. Math. **46**, 315–322 (1986).
- [G-VI] van der Geer, G.; van der Vlugt, M.: How to construct curves over finite fields with many points, alg-geom e-prints 9511005.
- [Go] Goppa, V.D.: Algebraic-Geometric codes, Math. USSR-Izv. **21** (1), 75–91 (1983).
- [H] Hansen, J.P.: Deligne-Lusztig varieties and group codes, Lect. Notes Math. **1518**, 63–81 (1992).
- [H-Sti] Hansen, J.P.; Stichtenoth, H.: Group codes on certain algebraic curves with many rational points, AAECC **1**, 67–77 (1990).
- [Har] Hartshorne, R.: Algebraic Geometry, Grad. Texts in Math. **52** Springer-Verlag 1977.
- [H-V] Hefez, A.; Voloch, J.F.: Frobenius non-classical curves, Arch. Math. **54**, 263–273 (1990).
- [Ho] Homma, M.: Linear systems on curves with no Weierstrass points, Bol. Soc. Bras. Mat. (N.S.) **23**, (1–2), 93–108 (1992).
- [I] Ibukiyama, T.: On rational points of curves of genus 3 over finite fields, Tôhoku Math. J. **45**, 311–329 (1993).
- [Ih] Ihara, Y.: Some Remarks on the Number of Rational Points of Algebraic Curves over Finite Fields, J. Fac. Sci. Tokyo **28**, 721–724 (1981).
- [L] Lang, S.: Abelian Varieties, Interscience Pub., New York 1959.
- [Le] Lewittes, J.: Places of degree one in function fields over finite fields, J. Pure Appl. Alg. **69**, 177–183 (1990).
- [P] Pedersen, J.P.: A function field related to the Ree group, Lect. Notes Math. **1518**, 122–131 (1992).
- [Ra] Rathmann, J.: The uniform position principle for curves in characteristic p , Math. Ann. **276**, 565–579 (1987).
- [R-Sti] Rück, H.G.; Stichtenoth, H.: A characterization of Hermitian function fields over finite fields, J. Reine Angew. Math. **457**, 185–188 (1994).
- [Sch] Schmidt, F.K.: Zur arithmetischen Theorie der algebraischen Funktionen II. Allgemeine Theorie der Weierstrasspunkte, Math. Z. **45**, 75–96 (1939).
- [Se] Serre, J.P.: Résumé des cours de 1983–1984, Annuaire du College de France, 79–83 (1984). Reprinted in Œuvres III, 701–705.
- [Sti] Stichtenoth, H.: Algebraic functions fields and codes, Springer-Verlag, Berlin, 1993.

- [Sti-X] Stichtenoth, H.; Xing, C.: The genus of maximal functions fields, Manuscripta Math. **86**, 217–224 (1995).
- [S-V] Stöhr, K.O.; Voloch, J.F.: Weierstrass points and curves over finite fields, Proc. London Math. Soc. (3) **52**, 1–19 (1986).
- [Ta] Tate, J.: Endomorphisms of abelian varieties over finite fields, Inventiones Math. **2**, 134–144 (1966).
- [To] Thomas, M.: Über das Geschlecht eines Funktionenkörpers, Diplomarbeit Universität Essen, 1994.

FACHBEREICH 6, MATHEMATIK UND INFORMATIK, UNIVERSITÄT ESSEN, D-45117, ESSEN - GERMANY

E-mail address: `RAINER.FUHRMANN@ZENTRALE.DEUTSCHE-BANK.dbp.de`

IMPA, ESTRADA DONA CASTORINA 110, 22460-320, RIO DE JANEIRO - BRAZIL

E-mail address: `garcia@impa.br`

FACHBEREICH 6, MATHEMATIK UND INFORMATIK, UNIVERSITÄT ESSEN, D-45117, ESSEN - GERMANY

E-mail address: `fernando.torres@uni-essen.de`